

Reliability and Validity of Borderline Personality Disorder in Hospitalized Adolescents

Catherine R. Glenn PhD¹; E. David Klonsky PhD²

Abstract

Objective: Although the DSM-IV suggests that dysfunctional personality patterns can be traced back to adolescence, there is continued debate about whether borderline personality disorder (BPD) can be reliably and validly diagnosed before age 18. The current study examined the reliability and validity of BPD in a large sample of adolescent psychiatric patients.

Method: BPD and Axis I disorders were assessed with validated structured interviews and a series of clinical, emotion, and personality correlates were assessed with validated self-report questionnaires. **Results:** Consistent with previous studies in adolescent clinical samples, approximately 30% of patients in the current sample met criteria for BPD. The nine BPD criteria demonstrated good internal consistency, equivalent to rates reported in adult samples. In addition, BPD was related to greater clinical severity and impairment as indexed by strong associations with all major Axis I disorders, as well as with dimensional measures of depression, anxiety, difficulties with emotion regulation, and impulsiveness. Notably, reliability and validity remained satisfactory even when analyses were limited to younger adolescents between the ages of 12 and 14.

Conclusions: Overall, findings suggest that BPD can be reliably and validly diagnosed in adolescents as young as 12-14 years old.

Key Words: *borderline personality disorder; adolescence; reliability; validity*

Résumé

Objectif: Bien que le DSM-IV suggère que les modèles de personnalité dysfonctionnelle puissent remonter à l'adolescence, un débat a encore cours à savoir si le trouble de la personnalité limite (TPL) peut être diagnostiqué avec fiabilité et validité avant l'âge de 18 ans. La présente étude a examiné la fiabilité et la validité du TPL dans un vaste échantillon d'adolescents patients psychiatriques. **Méthode:** Le TPL et les troubles de l'axe I ont été évalués à l'aide d'entrevues structurées validées, et une série de corrélats cliniques, émotionnels et de la personnalité ont été évalués par des questionnaires d'auto-évaluation validés. **Résultats:** Conformément aux études précédentes d'échantillons cliniques d'adolescents, environ 30% des patients du présent échantillon satisfaisaient aux critères du TPL. Les neuf critères du TPL démontraient une bonne cohésion interne, équivalente aux taux déclarés dans les échantillons d'adultes. En outre, le TPL était relié à une gravité et une incapacité cliniques accrues, comme l'indiquaient les fortes associations avec tous les troubles majeurs de l'axe I, ainsi qu'avec les mesures dimensionnelles d'éléments comme la dépression, l'anxiété, les difficultés de régulation émotionnelle, et l'impulsivité. Notamment, la fiabilité et la validité sont demeurées satisfaisantes même lorsque les analyses étaient limitées à de jeunes adolescents de 12 à 14 ans. **Conclusions:** Globalement, les résultats suggèrent que le TPL peut être diagnostiqué avec fiabilité et validité chez les adolescents dès l'âge de 12 à 14 ans.

Mots clés: *trouble de la personnalité limite; adolescence; fiabilité; validité*

Borderline personality disorder (BPD) is a debilitating clinical disorder characterized by significant impairment in affective, interpersonal, and behavioral domains (APA, 2000). In adults, rates of BPD are approximately

1-2% in the general population (Torgersen, Kringlen, & Cramer, 2001; Widiger & Weissman, 1991) and 15-50% in patient samples, depending on the severity of the clinical group (Becker, Grilo, Edell, & McGlashan, 2002; Widiger

¹Department of Psychology, Harvard University, Cambridge, Massachusetts, USA

²Department of Psychology, University of British Columbia, Vancouver, British Columbia

Corresponding E-Mail: catherineglenn@fas.harvard.edu

Submitted: April 3, 2013; Accepted: May 9, 2013

& Weissman, 1991). Current estimates suggest that rates of BPD in adolescent samples are similar, but somewhat higher, with 2-10% of adolescents in community samples (Bernstein et al., 1993; Leung & Leung, 2009) and 50% of adolescents in clinical samples meeting diagnostic criteria for BPD (Becker et al., 2002; Levy et al., 1999).

Although the DSM-IV suggests that dysfunctional personality patterns can be traced back to adolescence, debate still remains about whether personality disorders (PDs) can be reliably and validly diagnosed before age 18 (see review: Miller, Muehlenkamp, & Jacobson, 2008). The main argument against diagnosing PDs in adolescents has been that personality is unstable until adulthood and therefore dysfunction in personality cannot be reasonably assessed during the adolescent years. However, growing research refutes this central argument by indicating that: (1) personality is actually relatively stable during adolescence (McCrae et al., 2002); (2) personality *pathology*, and BPD in particular, is also moderately stable in adolescent samples (e.g., Bernstein et al., 1993; Mattanah, Becker, Levy, Edell, & McGlashan, 1995); and, (3) personality features underlying BPD (e.g., relational aggression) may be stable in children as young as 6-12 years of age (Stepp, Pilkonis, Hipwell, Loeber, & Stouthamer-Loeber, 2010).

Moreover, previous research also indicates that personality pathology among adolescents may be associated with significant impairment in this age group. For instance, BPD features in adolescents have been related to clinically significant behaviors, such as suicidal thoughts and behaviors (Glenn, Bagge, & Osman, in press), as well as have been found to prospectively predict impairment in interpersonal (Daley, Burge, & Hammen, 2000; Winograd, Cohen, & Chen, 2008) and academic/occupational domains (Winograd et al., 2008). Given the stability and potential clinical implications of personality pathology in adolescents, it is important for research to examine whether BPD diagnoses can be reliably and validly diagnosed in this age group.

Importantly, research has started to examine BPD pathology in adolescents and found that BPD diagnoses in this age group exhibit some similarities to adult BPD. For instance, the internal consistency of BPD criteria (Becker et al., 1999) and predictive utility of individual BPD criteria appear similar in adolescents and adults (Becker et al., 2002). Further, in line with adults, BPD pathology in adolescents has demonstrated reasonable stability over short (Bernstein et al., 1993; Mattanah et al., 1995) and long intervals (Winograd et al., 2008), as well as good predictive validity of clinical impairment indices (Levy et al., 1999; Winograd et al., 2008).

Although increasing research has examined BPD in younger age groups, few studies have examined how the reliability and validity of BPD may vary across adolescence. For instance, Bernstein et al. (1993) compared the prevalence and stability of all DSM-III-R PDs in early, middle,

and late adolescents, but not the reliability and validity of BPD diagnoses between these different age groups. Given that adolescence spans a wide range from post-childhood to emerging adulthood, it is important to examine whether the psychometric properties of BPD diagnoses change during this developmental period. Therefore, the purpose of the current study was to examine the reliability (i.e., internal consistency) and validity (i.e., predictive validity) of BPD in a large sample of adolescent psychiatric patients, and specifically to examine differences in BPD between younger (ages 12-14) and older (ages 15-18) adolescents.

Method

Participants and Procedure

Adolescents for the current study were recruited from the inpatient and partial hospitalization units of a hospital in the northeastern U.S. that provides short-term treatment for adolescents with severe psychopathology (e.g., range of major Axis I disorders, BPD, and self-injurious behaviors). The sample for the current study consisted of 174 adolescents (75.9% female) who completed the BPD interview measure (SIDP) described below. The largest ethnic groups were Caucasian (63.8%), Hispanic (12.6%), African American (10.9%), and mixed ethnic background (11.5%). Adolescents ranged in age from 12 (one participant) to 18 (one participant) (M age = 15.13; SD = 1.38). Because we were interested in comparing younger and older adolescents, participants were split into two groups: 12-14 years old (n = 62) and 15-18 years old (n = 112) that correspond to the middle school and high school years, respectively.

Detailed recruitment and data collection procedures for this project have been reported in a previous manuscript (see Glenn & Klonsky, in press), but will be summarized briefly here. IRB approval and informed consent/assent were obtained prior to study initiation. Participants were recruited for a larger study on nonsuicidal self-injury (NSSI) and therefore rates of NSSI are high in this sample. Adolescents completed study measures in one to two sessions at the hospital. A masters-level doctoral student, trained to reliability on measures of Axis I and II disorders (i.e., $r_s \geq .90$ with other masters- or doctoral-level trained interviewers) completed all interviews for the current project.

Interviews

Structured Interview for DSM-IV Personality (SIDP-IV)

The SIDP-IV (Pfohl, Blum, & Zimmerman, 1997) is a semi-structured interview that assesses all DSM-IV personality disorders including BPD. Each BPD criterion is rated on the following scale: 0 = *not present*, 1 = *subthreshold*, 2 = *present*, and 3 = *strongly present, associated with subjective distress*. Dimensional BPD scores are obtained by summing the 0-3 score for each criterion. A BPD criterion is considered present if rated as a 2 or 3. Importantly, the

Table 1. Borderline personality disorder (BPD) criteria endorsed

BPD criteria ($\alpha = .81$)	% endorsed in total sample (12-14 years / 15-18 years) ^a	Sensitivity %	Specificity %	Positive predictive value %	Negative predictive value %
Avoid abandonment	17.2 (12.9 / 19.6)	41.4	94.8	80.0	76.4
Unstable relationships	20.1 (21.0 / 19.6)	46.6	93.1	77.1	77.7
Identity disturbance	27.2 (24.2 / 28.8)	69.0	93.9	85.1	85.7
Impulsiveness	51.1 (33.9 / 60.7)	79.3	62.9	51.7	85.9
Suicidal/self-harm behaviors	58.6 (58.1 / 58.9)	94.8	59.5	53.9	95.8
Affective instability	47.1 (40.3 / 50.9)	86.2	72.4	61.0	91.3
Emptiness	39.1 (38.7 / 39.3)	75.9	79.3	64.7	86.8
Inappropriate anger	73.6 (69.4 / 75.9)	93.1	36.2	42.2	91.3
Dissociation/paranoia	27.6 (22.6 / 30.4)	62.1	89.7	75.0	82.5

^aPercentage of adolescents endorsing criterion as 2 or 3 on the Structured Interview for DSM-IV Personality Disorders in the total sample ($n=174$), as well as within the 12-14 year old and 15-18 year old subgroups.

SIDP-IV has demonstrated good psychometric properties in adolescents (Brent, Zelenak, Burstein, & Brown, 1990).

Mini-International Neuropsychiatric Interview for Children and Adolescents, English Version 6.0 (MINI-Kid).

The MINI-Kid (Sheehan, Shytle, Milo, Janavs, & Lecrubier, 2009) is a brief, structured interview that assesses all major DSM-IV Axis I disorders diagnosed during childhood and adolescence. The MINI-Kid has demonstrated good to excellent test-retest and interrater reliability, as well as good concordance with other clinical interviews for children (Sheehan et al., 2010).

Self-Report Questionnaires

Depression Anxiety Stress Scales (DASS-21)

The DASS-21 (Lovibond & Lovibond, 1995), a dimensional measure of depression, anxiety, and stress, has demonstrated good internal consistency and validity in previous research (Antony, Bieling, Cox, Enns, & Swinson, 1998).

Difficulties in Emotion Regulation Scale (DERS)

The DERS, a 36-item scale that assesses six different aspects of emotion regulation difficulties (see Gratz & Roemer, 2004), has demonstrated good reliability (internal

consistency and test-retest reliability) and construct validity in adolescent samples (Weinberg & Klonsky, 2009).

UPPS Impulsive Behavior Scale (UPPS)

The UPPS (Whiteside & Lynam, 2001), a 45-item measure of four distinct pathways to impulsive behavior, has been validated in previous studies (Whiteside, Lynam, Miller, & Reynolds, 2005).

Results

Descriptives and Reliability

Fifty-eight adolescents (33.3% of the total sample) met full criteria for a BPD diagnosis (M criteria endorsed = 6.48, $SD = 1.26$). Adolescents with BPD were somewhat older ($M = 15.47$, $SD = 1.43$) than adolescents without BPD ($M = 14.96$, $SD = 1.33$), $t(172) = 2.31$, $p = .022$. However, there was a non-significant difference between the number of adolescents with BPD in the 12-14 year-old group (29.0%) and the 15-18 year-old group (35.7%), $\chi^2(1, N = 174) = 0.80$, $p = .361$.

Table 1 displays the BPD criteria endorsed by adolescents in the total sample, as well as within the two age groups. Internal consistency of the BPD criteria was good in the total

Table 2. Prevalence of major Axis I disorders among adolescents with and without BPD

Axis I Disorder ^a	BPD (<i>n</i> = 58)	non-BPD (<i>n</i> = 116)	χ^2	<i>df</i>	Odds Ratio	95% CI
Alcohol/substance use	63.0%	30.0%	16.29***	1, 164	3.97	2.00-7.88
Anxiety	83.6%	45.0%	22.46***	1, 166	6.24	2.78-13.97
Behavioral	83.3%	63.1%	7.08**	1, 165	2.93	1.30-6.60
Mood	78.2%	35.4%	27.09***	1, 168	6.54	3.10-13.80

p* < .01, *p* < .001

^aAlcohol/Substance use disorder includes presence of current alcohol abuse/dependence or substance abuse/dependence. Anxiety disorder includes presence of any of the following current disorders: panic disorder, agoraphobia, social phobia, specific phobia, obsessive-compulsive disorder, posttraumatic stress disorder, or generalized anxiety disorder. Behavioral disorder includes presence of current attention-deficit hyperactivity disorder, conduct disorder or oppositional defiant disorder. Mood disorder includes presence of current bipolar I, bipolar II, major depressive disorder, or dysthymia.

Table 3. Bivariate correlations between BPD symptoms and clinical, emotion, and personality variables

Correlates (α)	Mean (SD)	BPD dimensional score		
		Total sample	12-14 years	15-18 years
Depression (.94)	9.37 (7.22)	.52***	.60***	.46***
Anxiety (.88)	8.00 (6.36)	.41***	.49***	.33**
Stress (.88)	9.50 (6.09)	.50***	.63***	.39***
Emotion regulation difficulties (.94)	104.65 (30.07)	.63***	.70***	.58***
Impulsiveness:				
Urgency (.84)	34.68 (7.46)	.48***	.53***	.42***
Lack of premeditation (.86)	26.82 (7.22)	.18*	.38**	.07
Lack of perseverance (.84)	25.26 (5.55)	.28**	.38**	.22*
Sensation seeking (.75)	31.34 (8.04)	.06	-.07	.13

p* < .05, *p* < .01, ****p* < .001

sample ($\alpha = .81$) and in the two age groups (12-14 year olds: $\alpha = .85$; 15-18 year olds: $\alpha = .78$). The only criterion-level difference in the two age groups was for impulsiveness, with greater endorsement in older compared to younger adolescents, $\chi^2(1, N = 174) = 11.51, p = .001$ (all other *ps* > .18). All item-total correlations were high (all *rs* > .30), except for impulsiveness ($r = .29$ in total sample), which was the only item that increased the internal consistency if deleted (to $\alpha = .82$ in total sample).

Next, we examined the predictive value of the individual BPD criteria. Suicidal/self-harm behaviors and inappropriate anger exhibited the greatest sensitivity in predicting presence of BPD, whereas unstable interpersonal relationships, efforts to avoid abandonment, and identity disturbance demonstrated the greatest specificity in predicting absence of BPD. Efforts to avoid abandonment and identity disturbance also exhibited the greatest positive predictive power, whereas suicidal/self-harm behaviors, inappropriate anger, and affective instability exhibited the greatest negative predictive power (see Table 1).

Validity

Next, we assessed the predictive validity of BPD in adolescents by examining associations with theoretically related constructs, including clinical, emotion, and personality correlates. Rates of all major Axis I disorders were significantly higher among adolescents with BPD, compared to those without BPD (see Table 2). Notably, this pattern was similar for the 12-14 year-old and 15-18 year-old groups, with one exception: among 15-18 year-olds, behavioral disorders were not significantly higher in the BPD group compared to the non-BPD group ($p = .130$), whereas this group difference was significant in 12-14 year-olds ($p = .021$).

Finally, we examined associations between BPD dimensional scale scores and a range of emotion and personality correlates (see Table 3). BPD symptoms were significantly related to depression, anxiety, and stress (DASS) scales, emotion regulation difficulties (DERS), and all impulsiveness scales (UPPS), except for Sensation Seeking ($p = .46$). Again, the pattern of associations was similar for the two age groups, although there was a non-significant tendency for correlations to be larger among younger adolescents (all *ps* > .06).

Discussion

Rates of BPD in the current sample were consistent with rates found in other adolescent psychiatric samples (e.g., Becker et al., 2002). The most frequently endorsed BPD criteria were inappropriate anger, suicidal behaviors, impulsiveness, and affective instability, and criteria endorsement was relatively similar for younger and older adolescents. The nine BPD criteria demonstrated good internal consistency in both younger and older adolescents, and these rates were consistent with those reported in adult samples (Becker et al., 1999). In addition, the item-total correlations for each of the BPD criteria were high in all adolescents, except for impulsiveness; removing the impulsiveness criterion improved the internal consistency of the remaining BPD criteria. This finding is consistent with Becker et al. (Becker, McGlashan, & Grilo, 2006) which found that impulsiveness was the only BPD criterion to load on its own factor; thus, for both adolescents and adults, impulsiveness may be a less useful indicator of BPD than other DSM-IV BPD criteria.

Findings also help clarify the predictive utility of the nine BPD criteria. Inappropriate anger and suicidal/self-harm behaviors provided the greatest sensitivity in predicting BPD diagnoses, whereas efforts to avoid abandonment, unstable relationships, and identity disturbance provided the greatest specificity. Identity disturbance also provided the highest positive predictive value for BPD diagnoses, whereas suicidal self-harm behaviors provided the highest negative predictive value. These results are generally consistent with previous studies (e.g., Becker et al., 2002; Pfohl, Coryell, Zimmerman, & Stangl, 1986), which have indicated that, although some criteria may provide more predictive value than others (e.g., identity disturbance and suicidal/self-harm behaviors), no single BPD criterion is pathognomonic of BPD. Instead, clusters or combinations of criteria are necessary to accurately distinguish adolescents with and without BPD.

Results from the current study also provide support for the construct validity of BPD in adolescents. Specifically, compared to adolescents without BPD, those with BPD were more likely to meet criteria for a range of DSM-IV Axis I disorders, including mood, anxiety, substance-related, and behavioral disorders. In addition, BPD criteria (on a continuous scale) were significantly associated with theoretically related constructs, such as depression, anxiety, difficulties with emotion regulation, and impulsiveness, which provides further concurrent validity for the diagnosis of BPD in adolescents. Notably, all patterns with clinical, emotion, and personality correlates were similar when younger and older adolescents were examined separately. In fact, BPD pathology exhibited even higher reliability and stronger associations with related clinical constructs in younger, compared to older, adolescents. Taken together, these findings suggest that BPD can be reasonably diagnosed in adolescents as young as 12-14 years of age.

This line of research has a number of clinical implications. First, consistent research now indicates that, similar to depression and anxiety, BPD is a disorder that can manifest across the lifespan. Therefore, given the growing evidence that BPD *can* be reliably and validly diagnosed in adolescents, BPD should be routinely assessed in adolescents to improve case conceptualization and treatment planning. This research also has implications for future editions of the DSM. Since the construct of BPD seems equally relevant to adolescents as for adults, the DSM should be revised to make BPD diagnoses more appropriate for adolescents (e.g., shorter time frames for symptoms and examples that may be more relevant for adolescent presentations of the criteria).

In sum, the current study provides additional support for diagnosing BPD in adolescents. However, there are some important limitations to the present research that warrant discussion. First, the current adolescent sample was from one hospital in the northeastern U.S. and was predominantly female and Caucasian. Future research should examine BPD in adolescents from diverse sociodemographic backgrounds. Second, the current study examined BPD in a clinically severe sample of adolescents that was oversampled for NSSI. Studies should also examine reliability and validity of BPD symptoms and diagnoses in community and less severe clinical samples. Third, although the overall sample was relatively large, the sample of adolescents in the younger age group was not large enough for certain follow-up analyses, such as the examination comparing BPD in younger and older adolescents. Finally, the current study used a cross-sectional design, which does not provide information about the stability of BPD diagnoses over time. In line with previous studies (e.g., Mattanah et al., 1995), future research should use prospective designs to examine the course of BPD across adolescence.

Acknowledgments/Conflicts of Interest

This project was supported in part by an NIH grant (MH08009601) awarded to E. David Klonsky, Ph.D. Special thanks to Kristie Golden, Ph.D., as well as to the staff and patients at South Oaks Hospital – The Long Island Home for all their support with this project.

References

- Antony, M. M., Bieling, P. J., Cox, B. J., Enns, M. W., & Swinson, R. P. (1998). Psychometric properties of the 42-item and 21-item versions of the Depression Anxiety Stress Scales in clinical groups and a community sample. *Psychological Assessment, 10*, 176-181.
- APA. (2000). *Diagnostic and statistical manual of mental disorders* (4th TR (text revision) ed.). Washington, DC: American Psychiatric Association.
- Becker, D. F., Grilo, C. M., Edell, W. S., & McGlashan, T. H. (2002). Diagnostic efficiency of borderline personality disorder criteria in hospitalized adolescents: Comparison with hospitalized adults. *American Journal of Psychiatry, 159*, 2042-2047.
- Becker, D. F., Grilo, C. M., Morey, L. C., Walker, M. L., Edell, W. S., & McGlashan, T. H. (1999). Applicability of personality disorder

- criteria to hospitalized adolescents: Evaluation of internal consistency and criterion overlap. *Journal of the American Academy of Child and Adolescent Psychiatry*, 38, 200-205.
- Becker, D. F., McGlashan, T. H., & Grilo, C. M. (2006). Exploratory factor analysis of borderline personality disorder criteria in hospitalized adolescents. *Comprehensive Psychiatry*, 99, 99-105.
- Bernstein, D. P., Cohen, P., Velez, C. N., Schwab-Stone, M., Siever, L. J., & Shinsato, L. (1993). Prevalence and stability of the DSM-III-R personality disorders in a community-based sample of adolescents. *American Journal of Psychiatry*, 150, 1237-1243.
- Brent, D. A., Zelenak, J. P., Burstein, O., & Brown, R. V. (1990). Reliability and validity of the structured interview for personality disorders in adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 29, 349-354.
- Daley, S. E., Burge, D., & Hammen, C. (2000). Borderline personality disorder symptoms as predictors of four-year romantic relationship dysfunction in young women: Addressing issues of specificity. *Journal of Abnormal Psychology*, 109, 451-460.
- Glenn, C. R., Bagge, C. L., & Osman, A. (in press). Unique associations between borderline personality disorder features and suicide ideation and attempts in adolescents. *Journal of Personality Disorders*.
- Glenn, C. R., & Klonsky, E. D. (in press). Non-suicidal self-injury disorder: An empirical investigation in adolescent psychiatric patients. *Journal of Clinical Child and Adolescent Psychiatry*.
- Gratz, K. L., & Roemer, L. (2004). Multidimensional assessment of emotion regulation and dysregulation: Development, factor structure, and initial validation of the difficulties in emotion regulation scale. *Journal of Psychopathology and Behavioral Assessment*, 26, 41-54.
- Levy, K. N., Becker, D. F., Grilo, C. M., Mattanah, J. J. F., Garnet, K. E., Quinlan, D. M.,...McGlashan, T. H. (1999). Concurrent and predictive validity of the personality disorder diagnosis in adolescent inpatients. *American Journal of Psychiatry*, 156, 1522-1528.
- Leung, S. W., & Leung, F. (2009). Construct validity and prevalence rate of borderline personality disorder among Chinese adolescents. *Journal of Personality Disorders*, 23, 494-513.
- Lovibond, S. H., & Lovibond, P. F. (1995). *Manual for the Depression Anxiety Stress Scales*, (2nd ed.). Sydney, Australia: Psychology Foundation of Australia.
- Mattanah, J. J. F., Becker, D. F., Levy, K. N., Edell, W. S., & McGlashan, T. H. (1995). Diagnostic stability in adolescents followed up 2 years after hospitalization. *American Journal of Psychiatry*, 152, 889-894.
- McCrae, R. R., Costa, P. T., Terracciano, A., Parker, W. D., Mills, C. J., De Fruyt, F., & Mervielde, I. (2002). Personality trait development from age 12 to age 18: Longitudinal, cross-sectional, and cross-cultural analyses. *Journal of Personality and Social Psychology*, 83, 1456-1468.
- Miller, A. L., Muehlenkamp, J. J., & Jacobson, C. M. (2008). Fact or fiction: Diagnosing borderline personality disorder in adolescents. *Clinical Psychology Review*, 28, 969-981.
- Pfohl, B., Blum, N., & Zimmerman, M. (1997). *Structured interview for DSM-IV personality*. Washington DC: American Psychiatric Press.
- Pfohl, B., Coryell, W., Zimmerman, M., & Stangl, D. (1986). DSM-III personality disorders: Diagnostic overlap and internal consistency of individual DSM-III criteria. *Comprehensive Psychiatry*, 27, 21-34.
- Sheehan, D., V., Sheehan, K. H., Shytle, R. D., Janavs, J., Bannon, Y., Rogers, J. E.,...Wilkinson, B. (2010). Reliability and validity of the MINI international neuropsychiatric interview for children and adolescents (MINI-KID). *Journal of Clinical Psychiatry*, 71, 313-326.
- Sheehan, D., Shytle, D., Milo, K., Janavs, J., & Lecrubier, Y. (2009). *MINI International Neuropsychiatric Interview for Children and Adolescents, English Version 6.0*. Tampa, FL: University of South Florida.
- Stepp, S. D., Pilkonis, P. A., Hipwell, A. E., Loeber, R., & Stouthamer-Loeber, M. (2010). Stability of borderline personality disorder features in girls. *Journal of Personality Disorders*, 24, 460-472.
- Torgersen, S., Kringlen, E., & Cramer, V. (2001). The prevalence of personality disorders in a community sample. *Archives of General Psychiatry*, 58, 590-596.
- Weinberg, A. & Klonsky, E. D. (2009). Measurement of emotion dysregulation in adolescents. *Psychological Assessment*, 21, 616-621.
- Whiteside, S. P., & Lynam, D. R. (2001). The five factor model and impulsivity: Using a structural model of personality to understand impulsivity. *Personality and Individual Differences*, 30, 669-689.
- Whiteside, S. P., Lynam, D. R., Miller, J. D., & Reynolds, S. K. (2005). Validation of the UPPS Impulsive Behavior Scale: A four-factor model of impulsivity. *European Journal of Personality*, 19, 559-574.
- Widiger, T. A., & Weissman, M. M. (1991). Epidemiology of borderline personality disorder. *Hospital & Community Psychiatry*, 42, 1015-1021.
- Winograd, G., Cohen, P., & Chen, H. (2008). Adolescent borderline symptoms in the community: Prognosis for functioning over 20 years. *Journal of Child Psychology and Psychiatry*, 49, 933-941.